

SMOKING-ATTRIBUTABLE DEATHS (WOMEN) 4,429
 SMOKING-ATTRIBUTABLE YEARS OF POTENTIAL LIFE LOST (WOMEN) 74,457

Status Indicators (Women)

| | White (Non-Hispanic) | Black (Non-Hispanic) | American Indian/Alaskan Native (Non-Hispanic) | Asian/Pacific Islander (Non-Hispanic) | Hispanic | U.S. Data | State Overall Data | State Grade | State Rank |
|--|----------------------|----------------------|---|---------------------------------------|----------|-----------|--------------------|-------------|------------|
| Prevalence: | | | | | | | | | |
| Current Smoking, Adults (%) | 21.0 | 23.2 | | 6.8 | 13.9 | 20.7 | 19.7 | U | 11 |
| Ages 18-24 | 26.4 | 15.2 | | | 12.8 | 24.0 | 22.4 | | |
| Ages 25-44 | 27.8 | 25.0 | | 2.5 | 17.9 | 24.4 | 24.0 | | |
| Ages 45-64 | 21.1 | 31.0 | | | 10.9 | 21.4 | 20.6 | | |
| Ages 65+ | 9.7 | 9.8 | | | 3.0 | 9.2 | 9.2 | | |
| Current Smoking, Grades 9-12 (%) | | | | | | 27.7 | 28.9 | | |
| Smoking During Pregnancy, All Ages (%) | | | | | | 12.2 | 9.7 | U | 14 |
| Ages 15-19 | | | | | | 18.1 | 16.5 | | |
| Cessation: | | | | | | | | | |
| Trying to Quit, Adults (%) | | | | | | 44.9 | 50.6 | F | 21 |
| Trying to Quit, Grades 9-12 (%) | | | | | | 61.4 | 50.2 | | |
| Receiving Smoking Cessation Advice by Physician (%) | | | | | | 61.0 | 61.6 | U | 25 |
| No Health Insurance (%) | 9.4 | 22.3 | | 17.8 | 36.1 | 17.0 | 15.4 | F | 28 |
| Second-Hand Smoke: | | | | | | | | | |
| Reporting Smoke Ban —At Work (%) | | | | | | 74.1 | 75.6 | F | 18 |
| —At Home (%) | | | | | | 64.5 | 64.7 | F | 21 |
| Smoking-Related Diseases: | | | | | | | | | |
| Lung Cancer Death Rate (per 100,000) | 42.5 | 41.8 | | 12.7 | 8.1 | 40.7 | 41.5 | F | 27 |
| Chronic Obstructive Pulmonary Disease Death Rate (per 100,000) | 89.2 | 72.3 | | | 30.9 | 103.5 | 86.3 | U | 7 |
| Coronary Heart Disease Death Rate (per 100,000) | 176.2 | 195.7 | | 56.9 | 71.2 | 164.6 | 176.6 | F | 41 |
| Stroke Death Rate (per 100,000) | 46.4 | 63.7 | | 27.2 | 22.3 | 61.5 | 48.2 | S- | 2 |

Grading Key for Status Indicators **S** Satisfactory **S-** Satisfactory Minus **U** Unsatisfactory **F** Fail

Policy Indicators

| | Strength of Policy |
|---|--------------------|
| Cessation: | |
| Telephone Quitline (1-866-NJ-STOPS) | |
| Pregnancy-Specific Counseling on Telephone Quitline | |
| Medicaid Smoking Cessation Coverage | |
| Medicaid Coverage of Pregnancy-Specific Counseling | |
| Private Insurance Smoking Cessation Coverage | |
| Second-Hand Smoke: | |
| Restrictions on Second-Hand Smoke | |
| Tax: | |
| Cigarette Excise Tax | |
| Funding: | |
| Tobacco Prevention Funding | |
| Youth Access: | |
| Tobacco Sales Rate to Minors | |
| Youth Access Restrictions | |

Grading Key for Policy Indicators Meets Policy Limited Policy Weak Policy Minimal/No Policy

Demographics

| | Number (Percent) | Number (Percent) |
|---|-------------------|---------------------|
| | New Jersey | U.S. |
| Total Population of Women | 4,358,967 (52.1%) | 143,584,500 (51.9%) |
| Women by Race | | |
| White (non-Hispanic) | 2,944,748 (67.6%) | 99,298,461 (69.2%) |
| Black (non-Hispanic) | 626,465 (14.4%) | 18,537,763 (12.9%) |
| American Indian/Alaskan Native (non-Hispanic) | 18,817 (0.4%) | 1,441,671 (1.0%) |
| Asian/Pacific Islander (non-Hispanic) | 265,683 (6.1%) | 6,324,727 (4.4%) |
| Hispanic | 503,255 (11.5%) | 17,981,877 (12.5%) |
| Women by Age | | |
| 0-17 | 971,813 (22.3%) | 35,375,812 (24.6%) |
| 18-24 | 356,621 (8.2%) | 13,489,103 (9.4%) |
| 25-44 | 1,330,924 (30.5%) | 42,030,761 (29.3%) |
| 45-64 | 1,023,574 (23.5%) | 33,196,167 (23.1%) |
| 65+ | 676,034 (15.5%) | 19,492,657 (13.6%) |
| Lesbian-Headed Households | 8,347 (0.27%) | 293,365 (0.28%) |
| Median Earnings for Women (\$) | \$31,000 | \$28,000 |
| Women Residing in—Urban Areas | 4,093,361 (94.5%) | 113,984,742 (79.5%) |
| —Rural Areas | 238,176 (5.5%) | 29,383,601 (20.5%) |
| Women (25 or older) by Education: | | |
| Less than 12 years | 408,712 (13.5%) | 14,877,627 (15.7%) |
| High School Graduate | 1,123,260 (37.1%) | 31,412,377 (33.2%) |
| Some College or Associate Degree | 645,767 (21.3%) | 25,038,018 (26.4%) |
| Bachelor's Degree | 586,837 (19.4%) | 16,098,968 (17.0%) |
| Graduate Degree | 265,957 (8.8%) | 7,292,595 (7.7%) |

SMOKING-ATTRIBUTABLE DEATHS (WOMEN) 810
 SMOKING-ATTRIBUTABLE YEARS OF POTENTIAL LIFE LOST (WOMEN) 13,622

Status Indicators (Women)

| | White (Non-Hispanic) | Black (Non-Hispanic) | American Indian/Alaskan Native (Non-Hispanic) | Asian/Pacific Islander (Non-Hispanic) | Hispanic | U.S. Data | State Overall Data | State Grade | State Rank |
|--|----------------------|----------------------|---|---------------------------------------|----------|-----------|--------------------|-------------|------------|
| Prevalence: | | | | | | | | | |
| Current Smoking, Adults (%) | 22.6 | 20.0 | 10.1 | | 20.2 | 20.7 | 20.7 | F | 21 |
| Ages 18-24 | 31.7 | | | | 23.0 | 24.0 | 24.4 | | |
| Ages 25-44 | 28.8 | | 7.6 | | 21.0 | 24.4 | 23.4 | | |
| Ages 45-64 | 22.0 | | 14.9 | | 21.4 | 21.4 | 21.4 | | |
| Ages 65+ | 10.5 | | | | 12.5 | 9.2 | 11.1 | | |
| Current Smoking, Grades 9-12 (%) | | | | | | 27.7 | | | |
| Smoking During Pregnancy, All Ages (%) | | | | | | 12.2 | 9.8 | U | 15 |
| Ages 15-19 | | | | | | 18.1 | 12.5 | | |
| Cessation: | | | | | | | | | |
| Trying to Quit, Adults (%) | | | | | | 44.9 | 53.9 | F | 7 |
| Trying to Quit, Grades 9-12 (%) | | | | | | 61.4 | | | |
| Receiving Smoking Cessation Advice by Physician (%) | | | | | | 61.0 | 60.9 | U | 29 |
| No Health Insurance (%) | 21.5 | | 49.4 | | 30.7 | 17.0 | 29.5 | F | 51 |
| Second-Hand Smoke: | | | | | | | | | |
| Reporting Smoke Ban —At Work (%) | | | | | | 74.1 | 73.1 | F | 28 |
| —At Home (%) | | | | | | 64.5 | 65.8 | F | 14 |
| Smoking-Related Diseases: | | | | | | | | | |
| Lung Cancer Death Rate (per 100,000) | 33.1 | | | | 20.7 | 40.7 | 31.4 | U | 6 |
| Chronic Obstructive Pulmonary Disease Death Rate (per 100,000) | 128.2 | | 51.3 | | 68.4 | 103.5 | 121.9 | F | 39 |
| Coronary Heart Disease Death Rate (per 100,000) | 132.0 | 149.5 | 70.1 | | 113.7 | 164.6 | 129.1 | S- | 14 |
| Stroke Death Rate (per 100,000) | 55.2 | 51.3 | 50.2 | | 54.9 | 61.5 | 54.9 | U | 8 |

Grading Key for Status Indicators **S** Satisfactory **S-** Satisfactory Minus **U** Unsatisfactory **F** Fail

Policy Indicators

Strength of Policy

| | |
|---|--|
| Cessation: | |
| Telephone Quitline (1-800-4-CANCER) | |
| Pregnancy-Specific Counseling on Telephone Quitline | |
| Medicaid Smoking Cessation Coverage | |
| Medicaid Coverage of Pregnancy-Specific Counseling | |
| Private Insurance Smoking Cessation Coverage | |
| Second-Hand Smoke: | |
| Restrictions on Second-Hand Smoke | |
| Tax: | |
| Cigarette Excise Tax | |
| Funding: | |
| Tobacco Prevention Funding | |
| Youth Access: | |
| Tobacco Sales Rate to Minors | |
| Youth Access Restrictions | |

Grading Key for Policy Indicators Meets Policy Limited Policy Weak Policy Minimal/No Policy

Demographics

| | New Mexico | | U.S. | |
|---|------------|-----------|-------------|-----------|
| | Number | (Percent) | Number | (Percent) |
| Total Population of Women | 917,459 | (52.1%) | 143,584,500 | (51.9%) |
| Women by Race | | | | |
| White (non-Hispanic) | 406,815 | (44.3%) | 99,298,461 | (69.2%) |
| Black (non-Hispanic) | 13,185 | (1.4%) | 18,537,763 | (12.9%) |
| American Indian/Alaskan Native (non-Hispanic) | 101,315 | (11.0%) | 1,441,671 | (1.0%) |
| Asian/Pacific Islander (non-Hispanic) | 11,701 | (1.3%) | 6,324,727 | (4.4%) |
| Hispanic | 384,443 | (41.9%) | 17,981,877 | (12.5%) |
| Women by Age | | | | |
| 0-17 | 239,356 | (26.1%) | 35,375,812 | (24.6%) |
| 18-24 | 96,613 | (10.5%) | 13,489,103 | (9.4%) |
| 25-44 | 236,163 | (25.7%) | 42,030,761 | (29.3%) |
| 45-64 | 225,049 | (24.5%) | 33,196,167 | (23.1%) |
| 65+ | 120,278 | (13.1%) | 19,492,657 | (13.6%) |
| Lesbian-Headed Households | 2,595 | (0.38%) | 293,365 | (0.28%) |
| Median Earnings for Women (\$) | \$24,000 | | \$28,000 | |
| Women Residing in—Urban Areas | 698,570 | (75.5%) | 113,984,742 | (79.5%) |
| —Rural Areas | 226,159 | (24.5%) | 29,383,601 | (20.5%) |
| Women (25 or older) by Education: | | | | |
| Less than 12 years | 108,880 | (18.7%) | 14,877,627 | (15.7%) |
| High School Graduate | 185,042 | (31.8%) | 31,412,377 | (33.2%) |
| Some College or Associate Degree | 159,936 | (27.5%) | 25,038,018 | (26.4%) |
| Bachelor's Degree | 75,740 | (13.0%) | 16,098,968 | (17.0%) |
| Graduate Degree | 51,892 | (8.9%) | 7,292,595 | (7.7%) |

SMOKING-ATTRIBUTABLE DEATHS (WOMEN) 11,154
 SMOKING-ATTRIBUTABLE YEARS OF POTENTIAL LIFE LOST (WOMEN) 183,382

Status Indicators (Women)

| | White (Non-Hispanic) | Black (Non-Hispanic) | American Indian/ Alaskan Native (Non-Hispanic) | Asian/ Pacific Islander (Non-Hispanic) | Hispanic | U.S. Data | State Overall Data | State Grade | State Rank |
|--|-------------------------|-------------------------|--|---|----------|--------------|--------------------------|----------------|---------------|
| Prevalence: | | | | | | | | | |
| Current Smoking, Adults (%) | 22.6 | 17.7 | | 7.1 | 19.7 | 20.7 | 21.0 | F | 25 |
| Ages 18-24 | 39.0 | 27.8 | | | 26.9 | 24.0 | 33.3 | | |
| Ages 25-44 | 29.5 | 22.3 | | 8.4 | 21.3 | 24.4 | 25.9 | | |
| Ages 45-64 | 19.7 | 13.8 | | | 14.4 | 21.4 | 17.7 | | |
| Ages 65+ | 10.1 | 9.1 | | | 17.4 | 9.2 | 10.3 | | |
| Current Smoking, Grades 9-12 (%) | | | | | | 27.7 | 32.8 | | |
| Smoking During Pregnancy, All Ages (%) | | | | | | 12.2 | 9.0 | U | 9 |
| Ages 15-19 | | | | | | 18.1 | 15.7 | | |
| Cessation: | | | | | | | | | |
| Trying to Quit, Adults (%) | | | | | | 44.9 | 51.9 | F | 16 |
| Trying to Quit, Grades 9-12 (%) | | | | | | 61.4 | 59.6 | | |
| Receiving Smoking Cessation Advice by Physician (%) | | | | | | 61.0 | 64.4 | U | 15 |
| No Health Insurance (%) | 12.2 | 23.7 | | 33.9 | 31.1 | 17.0 | 18.8 | F | 42 |
| Second-Hand Smoke: | | | | | | | | | |
| Reporting Smoke Ban —At Work (%) | | | | | | 74.1 | 77.0 | F | 13 |
| —At Home (%) | | | | | | 64.5 | 62.9 | F | 28 |
| Smoking-Related Diseases: | | | | | | | | | |
| Lung Cancer Death Rate (per 100,000) | 40.2 | 31.0 | | 17.0 | 12.2 | 40.7 | 38.1 | F | 14 |
| Chronic Obstructive Pulmonary Disease Death Rate (per 100,000) | 92.1 | 50.0 | | 21.8 | 38.5 | 103.5 | 85.3 | S- | 6 |
| Coronary Heart Disease Death Rate (per 100,000) | 217.8 | 249.4 | 67.8 | 126.7 | 146.6 | 164.6 | 220.7 | F | 51 |
| Stroke Death Rate (per 100,000) | 42.0 | 41.6 | | 28.8 | 23.4 | 61.5 | 42.2 | S | 1 |

Grading Key for Status Indicators **S** Satisfactory **S-** Satisfactory Minus **U** Unsatisfactory **F** Fail

Policy Indicators

| | Strength of Policy |
|---|--------------------|
| Cessation: | |
| Telephone Quitline (1-866-NY-QUIT5) | |
| Pregnancy-Specific Counseling on Telephone Quitline | |
| Medicaid Smoking Cessation Coverage | |
| Medicaid Coverage of Pregnancy-Specific Counseling | |
| Private Insurance Smoking Cessation Coverage | |
| Second-Hand Smoke: | |
| Restrictions on Second-Hand Smoke | |
| Tax: | |
| Cigarette Excise Tax | |
| Funding: | |
| Tobacco Prevention Funding | |
| Youth Access: | |
| Tobacco Sales Rate to Minors | |
| Youth Access Restrictions | |

Grading Key for Policy Indicators Meets Policy Limited Policy Weak Policy Minimal/No Policy

Demographics

| | Number (Percent) | Number (Percent) |
|---|-------------------|---------------------|
| | New York | U.S. |
| Total Population of Women | 9,791,070 (52.9%) | 143,584,500 (51.9%) |
| Women by Race | | |
| White (non-Hispanic) | 6,250,989 (63.8%) | 99,298,461 (69.2%) |
| Black (non-Hispanic) | 1,551,664 (15.8%) | 18,537,763 (12.9%) |
| American Indian/Alaskan Native (non-Hispanic) | 29,232 (0.3%) | 1,441,671 (1.0%) |
| Asian/Pacific Islander (non-Hispanic) | 589,979 (6.0%) | 6,324,727 (4.4%) |
| Hispanic | 1,369,206 (14.0%) | 17,981,877 (12.5%) |
| Women by Age | | |
| 0-17 | 2,271,730 (23.2%) | 35,375,812 (24.6%) |
| 18-24 | 918,136 (9.4%) | 13,489,103 (9.4%) |
| 25-44 | 2,852,743 (29.1%) | 42,030,761 (29.3%) |
| 45-64 | 2,308,421 (23.6%) | 33,196,167 (23.1%) |
| 65+ | 1,440,040 (14.7%) | 19,492,657 (13.6%) |
| Lesbian-Headed Households | 21,996 (0.31%) | 293,365 (0.28%) |
| Median Earnings for Women (\$) | \$30,000 | \$28,000 |
| Women Residing in—Urban Areas | 8,652,632 (88.0%) | 113,984,742 (79.5%) |
| —Rural Areas | 1,177,077 (12.0%) | 29,383,601 (20.5%) |
| Women (25 or older) by Education: | | |
| Less than 12 years | 1,131,238 (17.1%) | 14,877,627 (15.7%) |
| High School Graduate | 2,196,039 (33.3%) | 31,412,377 (33.2%) |
| Some College or Associate Degree | 1,475,529 (22.4%) | 25,038,018 (26.4%) |
| Bachelor's Degree | 1,123,354 (17.0%) | 16,098,968 (17.0%) |
| Graduate Degree | 675,045 (10.2%) | 7,292,595 (7.7%) |

SMOKING-ATTRIBUTABLE DEATHS (WOMEN) 4,147
 SMOKING-ATTRIBUTABLE YEARS OF POTENTIAL LIFE LOST (WOMEN) 76,720

Status Indicators (Women)

| | White (Non-Hispanic) | Black (Non-Hispanic) | American Indian/Alaskan Native (Non-Hispanic) | Asian/Pacific Islander (Non-Hispanic) | Hispanic | U.S. Data | State Overall Data | State Grade | State Rank |
|--|----------------------|----------------------|---|---------------------------------------|----------|-----------|--------------------|-------------|------------|
| Prevalence: | | | | | | | | | |
| Current Smoking, Adults (%) | 24.1 | 21.3 | 22.2 | 16.4 | 25.9 | 20.7 | 23.4 | F | 39 |
| Ages 18-24 | 36.7 | 22.3 | | | | 24.0 | 31.7 | | |
| Ages 25-44 | 28.9 | 26.2 | | | 22.7 | 24.4 | 27.6 | | |
| Ages 45-64 | 22.7 | 22.3 | | | | 21.4 | 22.9 | | |
| Ages 65+ | 11.9 | 5.8 | | | | 9.2 | 11.0 | | |
| Current Smoking, Grades 9-12 (%) | | | | | | 27.7 | 27.2 | | |
| Smoking During Pregnancy, All Ages (%) | | | | | | 12.2 | 14.0 | F | 29 |
| Ages 15-19 | | | | | | 18.1 | 18.5 | | |
| Cessation: | | | | | | | | | |
| Trying to Quit, Adults (%) | | | | | | 44.9 | 51.3 | F | 18 |
| Trying to Quit, Grades 9-12 (%) | | | | | | 61.4 | 63.1 | | |
| Receiving Smoking Cessation Advice by Physician (%) | | | | | | 61.0 | 64.1 | U | 16 |
| No Health Insurance (%) | 14.5 | 19.2 | | | 68.3 | 17.0 | 16.9 | F | 33 |
| Second-Hand Smoke: | | | | | | | | | |
| Reporting Smoke Ban —At Work (%) | | | | | | 74.1 | 65.8 | F | 49 |
| —At Home (%) | | | | | | 64.5 | 55.5 | F | 46 |
| Smoking-Related Diseases: | | | | | | | | | |
| Lung Cancer Death Rate (per 100,000) | 41.1 | 31.1 | 22.0 | | | 40.7 | 39.1 | F | 17 |
| Chronic Obstructive Pulmonary Disease Death Rate (per 100,000) | 112.3 | 51.8 | 68.7 | | | 103.5 | 101.5 | U | 23 |
| Coronary Heart Disease Death Rate (per 100,000) | 148.9 | 189.0 | 205.3 | 53.7 | 24.6 | 164.6 | 156.2 | U | 26 |
| Stroke Death Rate (per 100,000) | 72.0 | 95.2 | 65.2 | | | 61.5 | 76.1 | F | 47 |

Grading Key for Status Indicators **S** Satisfactory **S-** Satisfactory Minus **U** Unsatisfactory **F** Fail

Policy Indicators

| | Strength of Policy |
|---|----------------------|
| Cessation: | |
| Telephone Quitline | <input type="text"/> |
| Pregnancy-Specific Counseling on Telephone Quitline | <input type="text"/> |
| Medicaid Smoking Cessation Coverage | <input type="text"/> |
| Medicaid Coverage of Pregnancy-Specific Counseling | <input type="text"/> |
| Private Insurance Smoking Cessation Coverage | <input type="text"/> |
| Second-Hand Smoke: | |
| Restrictions on Second-Hand Smoke | <input type="text"/> |
| Tax: | |
| Cigarette Excise Tax | <input type="text"/> |
| Funding: | |
| Tobacco Prevention Funding | <input type="text"/> |
| Youth Access: | |
| Tobacco Sales Rate to Minors | <input type="text"/> |
| Youth Access Restrictions | <input type="text"/> |

Grading Key for Policy Indicators Meets Policy Limited Policy Weak Policy Minimal/No Policy

Demographics

| | Number (Percent) | Number (Percent) |
|---|-------------------|---------------------|
| | North Carolina | U.S. |
| Total Population of Women | 4,111,555 (52.4%) | 143,584,500 (51.9%) |
| Women by Race | | |
| White (non-Hispanic) | 2,783,334 (67.7%) | 99,298,461 (69.2%) |
| Black (non-Hispanic) | 961,623 (23.4%) | 18,537,763 (12.9%) |
| American Indian/Alaskan Native (non-Hispanic) | 127,516 (3.1%) | 1,441,671 (1.0%) |
| Asian/Pacific Islander (non-Hispanic) | 75,639 (1.8%) | 6,324,727 (4.4%) |
| Hispanic | 163,444 (4.0%) | 17,981,877 (12.5%) |
| Women by Age | | |
| 0-17 | 973,936 (23.7%) | 35,375,812 (24.6%) |
| 18-24 | 428,043 (10.4%) | 13,489,103 (9.4%) |
| 25-44 | 1,183,215 (28.8%) | 42,030,761 (29.3%) |
| 45-64 | 973,356 (23.7%) | 33,196,167 (23.1%) |
| 65+ | 553,006 (13.5%) | 19,492,657 (13.6%) |
| Lesbian-Headed Households | 8,349 (0.27%) | 293,365 (0.28%) |
| Median Earnings for Women (\$) | \$25,000 | \$28,000 |
| Women Residing in—Urban Areas | 2,494,951 (60.8%) | 113,984,742 (79.5%) |
| —Rural Areas | 1,611,667 (39.2%) | 29,383,601 (20.5%) |
| Women (25 or older) by Education: | | |
| Less than 12 years | 519,478 (19.2%) | 14,877,627 (15.7%) |
| High School Graduate | 901,428 (33.3%) | 31,412,377 (33.2%) |
| Some College or Associate Degree | 718,071 (26.5%) | 25,038,018 (26.4%) |
| Bachelor's Degree | 407,467 (15.0%) | 16,098,968 (17.0%) |
| Graduate Degree | 163,133 (6.0%) | 7,292,595 (7.7%) |

National and State Report Cards

SMOKING-ATTRIBUTABLE DEATHS (WOMEN) **267**
 SMOKING-ATTRIBUTABLE YEARS OF POTENTIAL LIFE LOST (WOMEN) **4,545**

Status Indicators (Women)

| | White (Non-Hispanic) | Black (Non-Hispanic) | American Indian/Alaskan Native (Non-Hispanic) | Asian/Pacific Islander (Non-Hispanic) | Hispanic | U.S. Data | State Overall Data | State Grade | State Rank |
|--|----------------------|----------------------|---|---------------------------------------|----------|-----------|--------------------|-------------|------------|
| Prevalence: | | | | | | | | | |
| Current Smoking, Adults (%) | 19.5 | | 45.7 | | | 20.7 | 20.4 | F | 17 |
| Ages 18-24 | 34.1 | | | | | 24.0 | 35.9 | | |
| Ages 25-44 | 23.4 | | | | | 24.4 | 24.3 | | |
| Ages 45-64 | 18.7 | | | | | 21.4 | 19.5 | | |
| Ages 65+ | 5.9 | | | | | 9.2 | 5.9 | | |
| Current Smoking, Grades 9-12 (%) | | | | | | 27.7 | 35.5 | | |
| Smoking During Pregnancy, All Ages (%) | | | | | | 12.2 | 18.0 | F | 41 |
| Ages 15-19 | | | | | | 18.1 | 35.2 | | |
| Cessation: | | | | | | | | | |
| Trying to Quit, Adults (%) | | | | | | 44.9 | 44.6 | F | 42 |
| Trying to Quit, Grades 9-12 (%) | | | | | | 61.4 | 62.1 | | |
| Receiving Smoking Cessation Advice by Physician (%) | | | | | | 61.0 | 43.8 | F | 51 |
| No Health Insurance (%) | 10.1 | | 34.1 | | | 17.0 | 12.7 | U | 19 |
| Second-Hand Smoke: | | | | | | | | | |
| Reporting Smoke Ban —At Work (%) | | | | | | 74.1 | 71.4 | F | 34 |
| —At Home (%) | | | | | | 64.5 | 61.5 | F | 31 |
| Smoking-Related Diseases: | | | | | | | | | |
| Lung Cancer Death Rate (per 100,000) | 30.1 | | | | | 40.7 | 30.6 | U | 3 |
| Chronic Obstructive Pulmonary Disease Death Rate (per 100,000) | 70.2 | | | | | 103.5 | 72.6 | S- | 3 |
| Coronary Heart Disease Death Rate (per 100,000) | 118.8 | | 228.5 | | | 164.6 | 120.8 | S- | 10 |
| Stroke Death Rate (per 100,000) | 58.1 | | | | | 61.5 | 58.7 | U | 16 |

Grading Key for Status Indicators **S** Satisfactory **S-** Satisfactory Minus **U** Unsatisfactory **F** Fail

Policy Indicators

| | Strength of Policy |
|---|-------------------------------------|
| Cessation: | |
| Telephone Quitline | <input type="checkbox"/> |
| Pregnancy-Specific Counseling on Telephone Quitline | <input type="checkbox"/> |
| Medicaid Smoking Cessation Coverage | <input checked="" type="checkbox"/> |
| Medicaid Coverage of Pregnancy-Specific Counseling | <input checked="" type="checkbox"/> |
| Private Insurance Smoking Cessation Coverage | <input type="checkbox"/> |
| Second-Hand Smoke: | |
| Restrictions on Second-Hand Smoke | <input checked="" type="checkbox"/> |
| Tax: | |
| Cigarette Excise Tax | <input type="checkbox"/> |
| Funding: | |
| Tobacco Prevention Funding | <input checked="" type="checkbox"/> |
| Youth Access: | |
| Tobacco Sales Rate to Minors | <input checked="" type="checkbox"/> |
| Youth Access Restrictions | <input type="checkbox"/> |

Grading Key for Policy Indicators Meets Policy Limited Policy Weak Policy Minimal/No Policy

Demographics

| | Number (Percent) | Number (Percent) |
|---|------------------|---------------------|
| | North Dakota | U.S. |
| Total Population of Women | 311,467 (50.6%) | 143,584,500 (51.9%) |
| Women by Race | | |
| White (non-Hispanic) | 281,570 (90.4%) | 99,298,461 (69.2%) |
| Black (non-Hispanic) | 1,451 (0.5%) | 18,537,763 (12.9%) |
| American Indian/Alaskan Native (non-Hispanic) | 23,672 (7.6%) | 1,441,671 (1.0%) |
| Asian/Pacific Islander (non-Hispanic) | 2,408 (0.8%) | 6,324,727 (4.4%) |
| Hispanic | 2,367 (0.8%) | 17,981,877 (12.5%) |
| Women by Age | | |
| 0-17 | 66,477 (21.3%) | 35,375,812 (24.6%) |
| 18-24 | 36,055 (11.6%) | 13,489,103 (9.4%) |
| 25-44 | 87,899 (28.2%) | 42,030,761 (29.3%) |
| 45-64 | 71,804 (23.1%) | 33,196,167 (23.1%) |
| 65+ | 49,232 (15.8%) | 19,492,657 (13.6%) |
| Lesbian-Headed Households | 343 (0.13%) | 293,365 (0.28%) |
| Median Earnings for Women (\$) | \$24,000 | \$28,000 |
| Women Residing in—Urban Areas | 181,679 (56.5%) | 113,984,742 (79.5%) |
| —Rural Areas | 139,997 (43.5%) | 29,383,601 (20.5%) |
| Women (25 or older) by Education: | | |
| Less than 12 years | 22,908 (11.0%) | 14,877,627 (15.7%) |
| High School Graduate | 62,280 (29.8%) | 31,412,377 (33.2%) |
| Some College or Associate Degree | 69,445 (33.2%) | 25,038,018 (26.4%) |
| Bachelor's Degree | 43,705 (20.9%) | 16,098,968 (17.0%) |
| Graduate Degree | 10,597 (5.1%) | 7,292,595 (7.7%) |

SMOKING-ATTRIBUTABLE DEATHS (WOMEN) 7,899
 SMOKING-ATTRIBUTABLE YEARS OF POTENTIAL LIFE LOST (WOMEN) 136,883

Status Indicators (Women)

| | White (Non-Hispanic) | Black (Non-Hispanic) | American Indian/Alaskan Native (Non-Hispanic) | Asian/Pacific Islander (Non-Hispanic) | Hispanic | U.S. Data | State Overall Data | State Grade | State Rank |
|--|----------------------|----------------------|---|---------------------------------------|----------|-----------|--------------------|-------------|------------|
| Prevalence: | | | | | | | | | |
| Current Smoking, Adults (%) | 26.2 | 24.1 | | | 31.9 | 20.7 | 26.2 | F | 48 |
| Ages 18-24 | 37.8 | 23.5 | | | | 24.0 | 36.4 | | |
| Ages 25-44 | 33.2 | 24.6 | | | | 24.4 | 32.5 | | |
| Ages 45-64 | 23.7 | 31.0 | | | | 21.4 | 24.7 | | |
| Ages 65+ | 11.5 | 8.4 | | | | 9.2 | 11.4 | | |
| Current Smoking, Grades 9-12 (%) | | | | | | 27.7 | 33.7 | | |
| Smoking During Pregnancy, All Ages (%) | | | | | | 12.2 | 19.0 | F | 45 |
| Ages 15-19 | | | | | | 18.1 | 28.4 | | |
| Cessation: | | | | | | | | | |
| Trying to Quit, Adults (%) | | | | | | 44.9 | 46.4 | F | 38 |
| Trying to Quit, Grades 9-12 (%) | | | | | | 61.4 | | | |
| Receiving Smoking Cessation Advice by Physician (%) | | | | | | 61.0 | 65.7 | S- | 10 |
| No Health Insurance (%) | 11.5 | 19.3 | | | 28.0 | 17.0 | 13.5 | U | 23 |
| Second-Hand Smoke: | | | | | | | | | |
| Reporting Smoke Ban —At Work (%) | | | | | | 74.1 | 70.6 | F | 40 |
| —At Home (%) | | | | | | 64.5 | 53.4 | F | 49 |
| Smoking-Related Diseases: | | | | | | | | | |
| Lung Cancer Death Rate (per 100,000) | 43.7 | 52.2 | | 24.8 | 12.7 | 40.7 | 44.3 | F | 38 |
| Chronic Obstructive Pulmonary Disease Death Rate (per 100,000) | 116.5 | 78.8 | | | 39.5 | 103.5 | 113.1 | F | 34 |
| Coronary Heart Disease Death Rate (per 100,000) | 169.9 | 210.2 | | 51.8 | 91.3 | 164.6 | 173.2 | F | 39 |
| Stroke Death Rate (per 100,000) | 59.8 | 75.6 | | 30.3 | 35.8 | 61.5 | 61.2 | U | 23 |

Grading Key for Status Indicators **S** Satisfactory **S-** Satisfactory Minus **U** Unsatisfactory **F** Fail

Policy Indicators

| | Strength of Policy |
|---|-------------------------------------|
| Cessation: | |
| Telephone Quitline | <input type="checkbox"/> |
| Pregnancy-Specific Counseling on Telephone Quitline | <input type="checkbox"/> |
| Medicaid Smoking Cessation Coverage | <input checked="" type="checkbox"/> |
| Medicaid Coverage of Pregnancy-Specific Counseling | <input type="checkbox"/> |
| Private Insurance Smoking Cessation Coverage | <input type="checkbox"/> |
| Second-Hand Smoke: | |
| Restrictions on Second-Hand Smoke | <input checked="" type="checkbox"/> |
| Tax: | |
| Cigarette Excise Tax | <input checked="" type="checkbox"/> |
| Funding: | |
| Tobacco Prevention Funding | <input checked="" type="checkbox"/> |
| Youth Access: | |
| Tobacco Sales Rate to Minors | <input checked="" type="checkbox"/> |
| Youth Access Restrictions | <input type="checkbox"/> |

Grading Key for Policy Indicators Meets Policy Limited Policy Weak Policy Minimal/No Policy

Demographics

| | Number (Percent) | Number (Percent) |
|---|-------------------|---------------------|
| | Ohio | U.S. |
| Total Population of Women | 5,753,598 (52.4%) | 143,584,500 (51.9%) |
| Women by Race | | |
| White (non-Hispanic) | 4,828,362 (83.9%) | 99,298,461 (69.2%) |
| Black (non-Hispanic) | 732,094 (12.7%) | 18,537,763 (12.9%) |
| American Indian/Alaskan Native (non-Hispanic) | 16,587 (0.3%) | 1,441,671 (1.0%) |
| Asian/Pacific Islander (non-Hispanic) | 87,095 (1.5%) | 6,324,727 (4.4%) |
| Hispanic | 89,460 (1.6%) | 17,981,877 (12.5%) |
| Women by Age | | |
| 0-17 | 1,320,490 (23.0%) | 35,375,812 (24.6%) |
| 18-24 | 514,738 (8.9%) | 13,489,103 (9.4%) |
| 25-44 | 1,689,650 (29.4%) | 42,030,761 (29.3%) |
| 45-64 | 1,402,479 (24.4%) | 33,196,167 (23.1%) |
| 65+ | 826,241 (14.4%) | 19,492,657 (13.6%) |
| Lesbian-Headed Households | 9,671 (0.22%) | 293,365 (0.28%) |
| Median Earnings for Women (\$) | \$28,000 | \$28,000 |
| Women Residing in—Urban Areas | 4,558,907 (78.1%) | 113,984,742 (79.5%) |
| —Rural Areas | 1,281,971 (21.9%) | 29,383,601 (20.5%) |
| Women (25 or older) by Education: | | |
| Less than 12 years | 465,265 (11.9%) | 14,877,627 (15.7%) |
| High School Graduate | 1,557,270 (39.7%) | 31,412,377 (33.2%) |
| Some College or Associate Degree | 1,014,202 (25.9%) | 25,038,018 (26.4%) |
| Bachelor's Degree | 625,790 (16.0%) | 16,098,968 (17.0%) |
| Graduate Degree | 255,843 (6.5%) | 7,292,595 (7.7%) |

SMOKING-ATTRIBUTABLE DEATHS (WOMEN) 2,202
 SMOKING-ATTRIBUTABLE YEARS OF POTENTIAL LIFE LOST (WOMEN) 40,006

Status Indicators (Women)

| | White (Non-Hispanic) | Black (Non-Hispanic) | American Indian/Alaskan Native (Non-Hispanic) | Asian/Pacific Islander (Non-Hispanic) | Hispanic | U.S. Data | State Overall Data | State Grade | State Rank |
|--|----------------------|----------------------|---|---------------------------------------|----------|-----------|--------------------|-------------|------------|
| Prevalence: | | | | | | | | | |
| Current Smoking, Adults (%) | 25.0 | 19.9 | 31.3 | 12.8 | 16.8 | 20.7 | 24.5 | F | 45 |
| Ages 18-24 | 29.3 | | 33.2 | | | 24.0 | 28.2 | | |
| Ages 25-44 | 32.1 | 13.2 | 38.8 | | 15.0 | 24.4 | 29.7 | | |
| Ages 45-64 | 26.0 | 29.5 | 28.9 | | 23.5 | 21.4 | 26.4 | | |
| Ages 65+ | 11.4 | 12.9 | 11.2 | | | 9.2 | 11.3 | | |
| Current Smoking, Grades 9-12 (%) | | | | | | 27.7 | | | |
| Smoking During Pregnancy, All Ages (%) | | | | | | 12.2 | 17.3 | F | 37 |
| Ages 15-19 | | | | | | 18.1 | 23.8 | | |
| Cessation: | | | | | | | | | |
| Trying to Quit, Adults (%) | | | | | | 44.9 | 53.0 | F | 11 |
| Trying to Quit, Grades 9-12 (%) | | | | | | 61.4 | | | |
| Receiving Smoking Cessation Advice by Physician (%) | | | | | | 61.0 | 58.9 | F | 33 |
| No Health Insurance (%) | 19.6 | 21.8 | 33.8 | | 38.1 | 17.0 | 22.2 | F | 48 |
| Second-Hand Smoke: | | | | | | | | | |
| Reporting Smoke Ban —At Work (%) | | | | | | 74.1 | 73.7 | F | 27 |
| —At Home (%) | | | | | | 64.5 | 57.4 | F | 43 |
| Smoking-Related Diseases: | | | | | | | | | |
| Lung Cancer Death Rate (per 100,000) | 45.9 | 43.8 | 25.8 | | | 40.7 | 44.4 | F | 39 |
| Chronic Obstructive Pulmonary Disease Death Rate (per 100,000) | 121.7 | 60.7 | 44.8 | | | 103.5 | 113.8 | F | 35 |
| Coronary Heart Disease Death Rate (per 100,000) | 193.5 | 230.2 | 106.5 | 99.4 | 71.0 | 164.6 | 189.5 | F | 47 |
| Stroke Death Rate (per 100,000) | 70.3 | 87.8 | 35.0 | | 33.1 | 61.5 | 69.2 | F | 41 |

Grading Key for Status Indicators **S** Satisfactory **S-** Satisfactory Minus **U** Unsatisfactory **F** Fail

Policy Indicators

| | Strength of Policy |
|---|--------------------|
| Cessation: | |
| Telephone Quitline (1-866-PITCH-EM) | |
| Pregnancy-Specific Counseling on Telephone Quitline | |
| Medicaid Smoking Cessation Coverage | |
| Medicaid Coverage of Pregnancy-Specific Counseling | |
| Private Insurance Smoking Cessation Coverage | |
| Second-Hand Smoke: | |
| Restrictions on Second-Hand Smoke | |
| Tax: | |
| Cigarette Excise Tax | |
| Funding: | |
| Tobacco Prevention Funding | |
| Youth Access: | |
| Tobacco Sales Rate to Minors | |
| Youth Access Restrictions | |

Grading Key for Policy Indicators Meets Policy Limited Policy Weak Policy Minimal/No Policy

Demographics

| | Number (Percent) | Number (Percent) |
|---|-------------------|---------------------|
| | Oklahoma | U.S. |
| Total Population of Women | 1,706,786 (51.4%) | 143,584,500 (51.9%) |
| Women by Race | | |
| White (non-Hispanic) | 1,342,708 (78.7%) | 99,298,461 (69.2%) |
| Black (non-Hispanic) | 115,642 (6.8%) | 18,537,763 (12.9%) |
| American Indian/Alaskan Native (non-Hispanic) | 137,953 (8.1%) | 1,441,671 (1.0%) |
| Asian/Pacific Islander (non-Hispanic) | 24,716 (1.4%) | 6,324,727 (4.4%) |
| Hispanic | 85,766 (5.0%) | 17,981,877 (12.5%) |
| Women by Age | | |
| 0-17 | 417,056 (24.4%) | 35,375,812 (24.6%) |
| 18-24 | 172,785 (10.1%) | 13,489,103 (9.4%) |
| 25-44 | 453,027 (26.5%) | 42,030,761 (29.3%) |
| 45-64 | 404,222 (23.7%) | 33,196,167 (23.1%) |
| 65+ | 259,697 (15.2%) | 19,492,657 (13.6%) |
| Lesbian-Headed Households | 2,952 (0.22%) | 293,365 (0.28%) |
| Median Earnings for Women (\$) | \$23,000 | \$28,000 |
| Women Residing in—Urban Areas | 1,159,920 (66.1%) | 113,984,742 (79.5%) |
| —Rural Areas | 594,839 (33.9%) | 29,383,601 (20.5%) |
| Women (25 or older) by Education: | | |
| Less than 12 years | 165,744 (14.8%) | 14,877,627 (15.7%) |
| High School Graduate | 442,442 (39.6%) | 31,412,377 (33.2%) |
| Some College or Associate Degree | 320,025 (28.7%) | 25,038,018 (26.4%) |
| Bachelor's Degree | 126,829 (11.4%) | 16,098,968 (17.0%) |
| Graduate Degree | 61,906 (5.5%) | 7,292,595 (7.7%) |

SMOKING-ATTRIBUTABLE DEATHS (WOMEN) 1,995
 SMOKING-ATTRIBUTABLE YEARS OF POTENTIAL LIFE LOST (WOMEN) 33,404

Status Indicators (Women)

| | White (Non-Hispanic) | Black (Non-Hispanic) | American Indian/Alaskan Native (Non-Hispanic) | Asian/Pacific Islander (Non-Hispanic) | Hispanic | U.S. Data | State Overall Data | State Grade | State Rank |
|--|----------------------|----------------------|---|---------------------------------------|----------|-----------|--------------------|-------------|------------|
| Prevalence: | | | | | | | | | |
| Current Smoking, Adults (%) | 19.4 | | 46.5 | 14.1 | 12.7 | 20.7 | 19.7 | U | 11 |
| Ages 18-24 | 29.0 | | | | | 24.0 | 26.2 | | |
| Ages 25-44 | 24.6 | | | | 16.0 | 24.4 | 24.3 | | |
| Ages 45-64 | 18.2 | | | | 3.8 | 21.4 | 18.6 | | |
| Ages 65+ | 9.0 | | | | | 9.2 | 9.6 | | |
| Current Smoking, Grades 9-12 (%) | | | | | | 27.7 | | | |
| Smoking During Pregnancy, All Ages (%) | | | | | | 12.2 | 13.4 | U | 27 |
| Ages 15-19 | | | | | | 18.1 | 25.2 | | |
| Cessation: | | | | | | | | | |
| Trying to Quit, Adults (%) | | | | | | 44.9 | 53.2 | F | 10 |
| Trying to Quit, Grades 9-12 (%) | | | | | | 61.4 | | | |
| Receiving Smoking Cessation Advice by Physician (%) | | | | | | 61.0 | 65.5 | U | 12 |
| No Health Insurance (%) | 12.1 | | | 17.3 | 38.6 | 17.0 | 14.4 | U | 26 |
| Second-Hand Smoke: | | | | | | | | | |
| Reporting Smoke Ban —At Work (%) | | | | | | 74.1 | 73.9 | F | 26 |
| —At Home (%) | | | | | | 64.5 | 73.8 | U | 6 |
| Smoking-Related Diseases: | | | | | | | | | |
| Lung Cancer Death Rate (per 100,000) | 46.4 | | | | | 40.7 | 45.7 | F | 41 |
| Chronic Obstructive Pulmonary Disease Death Rate (per 100,000) | 125.4 | | | | | 103.5 | 123.2 | F | 40 |
| Coronary Heart Disease Death Rate (per 100,000) | 115.9 | 138.0 | 106.6 | 76.2 | 63.9 | 164.6 | 115.7 | S- | 7 |
| Stroke Death Rate (per 100,000) | 80.0 | 96.2 | | 75.1 | 29.8 | 61.5 | 80.1 | F | 49 |

Grading Key for Status Indicators **S** Satisfactory **S-** Satisfactory Minus **U** Unsatisfactory **F** Fail

Policy Indicators

| | Strength of Policy |
|---|-------------------------------------|
| Cessation: | |
| Telephone Quitline | <input type="checkbox"/> |
| Pregnancy-Specific Counseling on Telephone Quitline | <input type="checkbox"/> |
| Medicaid Smoking Cessation Coverage | <input checked="" type="checkbox"/> |
| Medicaid Coverage of Pregnancy-Specific Counseling | <input checked="" type="checkbox"/> |
| Private Insurance Smoking Cessation Coverage | <input type="checkbox"/> |
| Second-Hand Smoke: | |
| Restrictions on Second-Hand Smoke | <input checked="" type="checkbox"/> |
| Tax: | |
| Cigarette Excise Tax | <input checked="" type="checkbox"/> |
| Funding: | |
| Tobacco Prevention Funding | <input checked="" type="checkbox"/> |
| Youth Access: | |
| Tobacco Sales Rate to Minors | <input checked="" type="checkbox"/> |
| Youth Access Restrictions | <input type="checkbox"/> |

Grading Key for Policy Indicators Meets Policy Limited Policy Weak Policy Minimal/No Policy

Demographics

| | Number (Percent) | Number (Percent) |
|---|-------------------|---------------------|
| | Oregon | U.S. |
| Total Population of Women | 1,750,599 (51.6%) | 143,584,500 (51.9%) |
| Women by Race | | |
| White (non-Hispanic) | 1,470,945 (84.0%) | 99,298,461 (69.2%) |
| Black (non-Hispanic) | 29,243 (1.7%) | 18,537,763 (12.9%) |
| American Indian/Alaskan Native (non-Hispanic) | 28,325 (1.6%) | 1,441,671 (1.0%) |
| Asian/Pacific Islander (non-Hispanic) | 77,334 (4.4%) | 6,324,727 (4.4%) |
| Hispanic | 144,752 (8.3%) | 17,981,877 (12.5%) |
| Women by Age | | |
| 0-17 | 424,117 (24.2%) | 35,375,812 (24.6%) |
| 18-24 | 157,691 (9.0%) | 13,489,103 (9.4%) |
| 25-44 | 521,662 (29.8%) | 42,030,761 (29.3%) |
| 45-64 | 421,418 (24.1%) | 33,196,167 (23.1%) |
| 65+ | 225,711 (12.9%) | 19,492,657 (13.6%) |
| Lesbian-Headed Households | 5,086 (0.38%) | 293,365 (0.28%) |
| Median Earnings for Women (\$) | \$27,000 | \$28,000 |
| Women Residing in—Urban Areas | 1,366,693 (79.2%) | 113,984,742 (79.5%) |
| —Rural Areas | 358,156 (20.8%) | 29,383,601 (20.5%) |
| Women (25 or older) by Education: | | |
| Less than 12 years | 145,531 (12.5%) | 14,877,627 (15.7%) |
| High School Graduate | 345,175 (29.5%) | 31,412,377 (33.2%) |
| Some College or Associate Degree | 373,075 (31.9%) | 25,038,018 (26.4%) |
| Bachelor's Degree | 211,956 (18.1%) | 16,098,968 (17.0%) |
| Graduate Degree | 93,053 (8.0%) | 7,292,595 (7.7%) |

SMOKING-ATTRIBUTABLE DEATHS (WOMEN) 7,779
 SMOKING-ATTRIBUTABLE YEARS OF POTENTIAL LIFE LOST (WOMEN) 130,901

Status Indicators (Women)

| | White (Non-Hispanic) | Black (Non-Hispanic) | American Indian/Alaskan Native (Non-Hispanic) | Asian/Pacific Islander (Non-Hispanic) | Hispanic | U.S. Data | State Overall Data | State Grade | State Rank |
|--|----------------------|----------------------|---|---------------------------------------|----------|-----------|--------------------|-------------|------------|
| Prevalence: | | | | | | | | | |
| Current Smoking, Adults (%) | 22.0 | 29.6 | | 15.1 | 31.5 | 20.7 | 22.8 | F | 37 |
| Ages 18-24 | 32.8 | 14.5 | | | | 24.0 | 32.2 | | |
| Ages 25-44 | 28.0 | 31.1 | | | 27.6 | 24.4 | 27.9 | | |
| Ages 45-64 | 22.5 | 42.4 | | | | 21.4 | 24.4 | | |
| Ages 65+ | 8.9 | 11.6 | | | | 9.2 | 9.0 | | |
| Current Smoking, Grades 9-12 (%) | | | | | | 27.7 | 28.3 | | |
| Smoking During Pregnancy, All Ages (%) | | | | | | 12.2 | 16.7 | F | 35 |
| Ages 15-19 | | | | | | 18.1 | 26.3 | | |
| Cessation: | | | | | | | | | |
| Trying to Quit, Adults (%) | | | | | | 44.9 | 44.1 | F | 45 |
| Trying to Quit, Grades 9-12 (%) | | | | | | 61.4 | | | |
| Receiving Smoking Cessation Advice by Physician (%) | | | | | | 61.0 | 65.6 | U | 11 |
| No Health Insurance (%) | 9.2 | 19.4 | | | 19.9 | 17.0 | 10.8 | U | 10 |
| Second-Hand Smoke: | | | | | | | | | |
| Reporting Smoke Ban —At Work (%) | | | | | | 74.1 | 74.0 | F | 25 |
| —At Home (%) | | | | | | 64.5 | 59.9 | F | 34 |
| Smoking-Related Diseases: | | | | | | | | | |
| Lung Cancer Death Rate (per 100,000) | 39.2 | 57.3 | | 12.9 | 16.4 | 40.7 | 40.5 | F | 22 |
| Chronic Obstructive Pulmonary Disease Death Rate (per 100,000) | 93.2 | 77.3 | | | 45.0 | 103.5 | 91.9 | U | 13 |
| Coronary Heart Disease Death Rate (per 100,000) | 167.2 | 189.2 | | 63.7 | 94.3 | 164.6 | 168.7 | F | 34 |
| Stroke Death Rate (per 100,000) | 57.2 | 73.1 | | 39.2 | 43.4 | 61.5 | 58.4 | U | 13 |

Grading Key for Status Indicators **S** Satisfactory **S-** Satisfactory Minus **U** Unsatisfactory **F** Fail

Policy Indicators

| | Strength of Policy |
|---|--------------------|
| Cessation: | |
| Telephone Quitline (1-877-724-1090) | |
| Pregnancy-Specific Counseling on Telephone Quitline | |
| Medicaid Smoking Cessation Coverage | |
| Medicaid Coverage of Pregnancy-Specific Counseling | |
| Private Insurance Smoking Cessation Coverage | |
| Second-Hand Smoke: | |
| Restrictions on Second-Hand Smoke | |
| Tax: | |
| Cigarette Excise Tax | |
| Funding: | |
| Tobacco Prevention Funding | |
| Youth Access: | |
| Tobacco Sales Rate to Minors | |
| Youth Access Restrictions | |

Grading Key for Policy Indicators Meets Policy Limited Policy Weak Policy Minimal/No Policy

Demographics

| | Number (Percent) | Number (Percent) |
|---|-------------------|---------------------|
| | Pennsylvania | U.S. |
| Total Population of Women | 6,254,395 (52.4%) | 143,584,500 (51.9%) |
| Women by Race | | |
| White (non-Hispanic) | 5,295,613 (84.7%) | 99,298,461 (69.2%) |
| Black (non-Hispanic) | 609,799 (9.7%) | 18,537,763 (12.9%) |
| American Indian/Alaskan Native (non-Hispanic) | 19,254 (0.3%) | 1,441,671 (1.0%) |
| Asian/Pacific Islander (non-Hispanic) | 107,828 (1.7%) | 6,324,727 (4.4%) |
| Hispanic | 221,900 (3.5%) | 17,981,877 (12.5%) |
| Women by Age | | |
| 0-17 | 1,384,403 (22.1%) | 35,375,812 (24.6%) |
| 18-24 | 537,784 (8.6%) | 13,489,103 (9.4%) |
| 25-44 | 1,758,799 (28.1%) | 42,030,761 (29.3%) |
| 45-64 | 1,616,841 (25.9%) | 33,196,167 (23.1%) |
| 65+ | 956,568 (15.3%) | 19,492,657 (13.6%) |
| Lesbian-Headed Households | 10,674 (0.22%) | 293,365 (0.28%) |
| Median Earnings for Women (\$) | \$29,000 | \$28,000 |
| Women Residing in—Urban Areas | 4,947,002 (77.9%) | 113,984,742 (79.5%) |
| —Rural Areas | 1,404,389 (22.1%) | 29,383,601 (20.5%) |
| Women (25 or older) by Education: | | |
| Less than 12 years | 625,878 (14.4%) | 14,877,627 (15.7%) |
| High School Graduate | 1,772,276 (40.9%) | 31,412,377 (33.2%) |
| Some College or Associate Degree | 914,580 (21.1%) | 25,038,018 (26.4%) |
| Bachelor's Degree | 660,877 (15.3%) | 16,098,968 (17.0%) |
| Graduate Degree | 358,597 (8.3%) | 7,292,595 (7.7%) |

SMOKING-ATTRIBUTABLE DEATHS (WOMEN) 770
 SMOKING-ATTRIBUTABLE YEARS OF POTENTIAL LIFE LOST (WOMEN) 12,008

Status Indicators (Women)

| | White (Non-Hispanic) | Black (Non-Hispanic) | American Indian/Alaskan Native (Non-Hispanic) | Asian/Pacific Islander (Non-Hispanic) | Hispanic | U.S. Data | State Overall Data | State Grade | State Rank |
|--|----------------------|----------------------|---|---------------------------------------|----------|-----------|--------------------|-------------|------------|
| Prevalence: | | | | | | | | | |
| Current Smoking, Adults (%) | 22.8 | 22.5 | | 14.3 | 15.8 | 20.7 | 22.3 | F | 35 |
| Ages 18-24 | 34.9 | | | | 14.6 | 24.0 | 29.6 | | |
| Ages 25-44 | 27.9 | 22.5 | | | 21.7 | 24.4 | 27.0 | | |
| Ages 45-64 | 22.6 | 31.9 | | | 6.0 | 21.4 | 22.2 | | |
| Ages 65+ | 11.6 | | | | | 9.2 | 11.3 | | |
| Current Smoking, Grades 9-12 (%) | | | | | | 27.7 | 25.6 | | |
| Smoking During Pregnancy, All Ages (%) | | | | | | 12.2 | 14.6 | F | 30 |
| Ages 15-19 | | | | | | 18.1 | 22.5 | | |
| Cessation: | | | | | | | | | |
| Trying to Quit, Adults (%) | | | | | | 44.9 | 44.0 | F | 47 |
| Trying to Quit, Grades 9-12 (%) | | | | | | 61.4 | 61.5 | | |
| Receiving Smoking Cessation Advice by Physician (%) | | | | | | 61.0 | 75.7 | S | 1 |
| No Health Insurance (%) | 6.7 | 16.1 | | | 24.0 | 17.0 | 8.7 | S- | 3 |
| Second-Hand Smoke: | | | | | | | | | |
| Reporting Smoke Ban —At Work (%) | | | | | | 74.1 | 74.0 | F | 22 |
| —At Home (%) | | | | | | 64.5 | 65.2 | F | 19 |
| Smoking-Related Diseases: | | | | | | | | | |
| Lung Cancer Death Rate (per 100,000) | 46.6 | | | | | 40.7 | 45.7 | F | 41 |
| Chronic Obstructive Pulmonary Disease Death Rate (per 100,000) | 94.3 | | | | | 103.5 | 93.4 | U | 16 |
| Coronary Heart Disease Death Rate (per 100,000) | 178.1 | 240.3 | | | 57.4 | 164.6 | 179.0 | F | 43 |
| Stroke Death Rate (per 100,000) | 50.6 | 94.3 | | | | 61.5 | 51.4 | S- | 6 |

Grading Key for Status Indicators **S** Satisfactory **S-** Satisfactory Minus **U** Unsatisfactory **F** Fail

Policy Indicators

| | Strength of Policy |
|---|--------------------|
| Cessation: | |
| Telephone Quitline (1-800-TRY-TO-STOP) | |
| Pregnancy-Specific Counseling on Telephone Quitline | |
| Medicaid Smoking Cessation Coverage | |
| Medicaid Coverage of Pregnancy-Specific Counseling | |
| Private Insurance Smoking Cessation Coverage | |
| Second-Hand Smoke: | |
| Restrictions on Second-Hand Smoke | |
| Tax: | |
| Cigarette Excise Tax | |
| Funding: | |
| Tobacco Prevention Funding | |
| Youth Access: | |
| Tobacco Sales Rate to Minors | |
| Youth Access Restrictions | |

Grading Key for Policy Indicators Meets Policy Limited Policy Weak Policy Minimal/No Policy

Demographics

| | Number (Percent) | Number (Percent) |
|---|------------------|---------------------|
| | Rhode Island | U.S. |
| Total Population of Women | 540,820 (53.8%) | 143,584,500 (51.9%) |
| Women by Race | | |
| White (non-Hispanic) | 459,017 (84.9%) | 99,298,461 (69.2%) |
| Black (non-Hispanic) | 20,328 (3.8%) | 18,537,763 (12.9%) |
| American Indian/Alaskan Native (non-Hispanic) | 4,142 (0.8%) | 1,441,671 (1.0%) |
| Asian/Pacific Islander (non-Hispanic) | 12,208 (2.3%) | 6,324,727 (4.4%) |
| Hispanic | 45,125 (8.3%) | 17,981,877 (12.5%) |
| Women by Age | | |
| 0-17 | 113,279 (20.9%) | 35,375,812 (24.6%) |
| 18-24 | 47,567 (8.8%) | 13,489,103 (9.4%) |
| 25-44 | 153,062 (28.3%) | 42,030,761 (29.3%) |
| 45-64 | 126,363 (23.4%) | 33,196,167 (23.1%) |
| 65+ | 100,549 (18.6%) | 19,492,657 (13.6%) |
| Lesbian-Headed Households | 1,299 (0.32%) | 293,365 (0.28%) |
| Median Earnings for Women (\$) | \$30,000 | \$28,000 |
| Women Residing in—Urban Areas | 496,739 (91.2%) | 113,984,742 (79.5%) |
| —Rural Areas | 47,945 (8.8%) | 29,383,601 (20.5%) |
| Women (25 or older) by Education: | | |
| Less than 12 years | 82,914 (21.8%) | 14,877,627 (15.7%) |
| High School Graduate | 110,296 (29.0%) | 31,412,377 (33.2%) |
| Some College or Associate Degree | 87,329 (23.0%) | 25,038,018 (26.4%) |
| Bachelor's Degree | 60,476 (15.9%) | 16,098,968 (17.0%) |
| Graduate Degree | 38,959 (10.3%) | 7,292,595 (7.7%) |

National and State Report Cards